

AD-A262 161



REPORT DOCUMENTATION PAGE

Form Approved OMB No. 0704-0188	
<p>Public reporting burden is estimated to average 1 hour per response. Subject to change. Comments concerning this burden estimate or any other aspect of the collection of information should be directed to the Office of Management and Budget, Paperwork Reduction Project (0704-0188), Executive Office of the President, Washington, DC 20503. On or before June 1, 1995.</p>	
1. AGENCY USE ONLY (Leave blank)	2. REPORT DATE
Reinterpreted	December 195 3
4. TITLE AND SUBTITLE	3. REPORT TYPE AND DATES COVERED
Resonance Electron Scattering by O ₂ Monolayers on Graphite: Reinterpreted	Technical
5. FUNDING NUMBERS	6. AUTHOR(S)
	Hideo Sabe and David E. Ramaker
7. PERFORMING ORGANIZATION NAME(S) AND ADDRESS(ES)	8. SPONSORING/MONITORING AGENCY NAME(S) AND ADDRESS(ES)
George Washington University Chemistry Department Washington, D.C. 20052	9. SPONSORING/MONITORING AGENCY NUMBER ARSENIC REPORT NUMBER
9. SPONSORING/MONITORING AGENCY NAME(S) AND ADDRESS(ES)	10. DISTRIBUTION CODE
Office of Naval Research 800 N. Quincy Street Arlington, VA 22217-5000	
11. SUPPLEMENTARY NOTES	12. DISTRIBUTION STATEMENT
Prepared for publication in Proc. of DIET V	Approved for public release; distribution is unlimited.
13. ABSTRACT (maximum 200 words)	
<p>Electron angular distributions of inelastic electron scattering from O₂ physisorbed on graphite have been studied extensively by Palser and co-workers. Their results have been published in more than eight papers over the past four years and summarized in two recent review articles. We have reinterpreted their data and some aspects come to different conclusions.</p>	
14. SUBJECT TERMS	15. NUMBER OF PAGES
Inelastic electron scattering, molecular oxygen, resonant states.	14. PRICE CODE
16. SECURITY CLASSIFICATION OF REPORT	17. SECURITY CLASSIFICATION OF THIS PAGE
Unclassified	Unclassified
Standard Form 295 (Rev. 7-89) GSA GEN. REG. NO. 27-14 MIL-STD-136A	

OFFICE OF NAVAL RESEARCH
CONTRACT NO0014-91-J-1896

RAT CODE 4134043-01

Technical Report No. 63

RESONANCE ELECTRON SCATTERING BY O₂
MONOLAYERS ON GRAPHITE: REINTERPRETED

BY

Hideo Sabe and David E. Ramaker

Prepared for Publication:

In the

Proc. of DIET V

DTIC
DRAFT
Sb
MAR 18 1993

George Washington University
Department of Chemistry
Washington, D.C. 20052

January 1993

Reproduction in whole or in part is permitted for
any purpose of the United States Government

- This document has been approved for public release
and sale; its distribution is unlimited.

93 3 17 044

93-05575



Resonance Electron Scattering by O₂ Monolayers on Graphite: Reinterpreted

Hideo Saito and David E. Palmer

Chemistry Department, George Washington University
Washington, D.C. 20052, USA

1. Introduction

Both electron stimulated desorption (ESD) and inelastic electron scattering (IES) are enhanced strongly by anion resonant states. Occasionally, peak features observed in ESD and IES arise from the same anion resonant states. IED, however, unlike ESD, provides information on the symmetry of the anion resonant states through the angular distributions of the inelastically scattered electrons. Such electron angular distributions from O₂ on graphite have been studied extensively by Palmer and co-workers. Their results have been published in more than eight papers over the past four years [1-8] and summarized in two recent review articles [7,8].

On graphite, O₂ can be prepared in either the 6 or f phase, where the molecules lie either parallel (6 phase) or perpendicular (f phase) to the graphite surface [9]. The excitation profile of the X(²P)⁻a(^3P)₀ electronic transition exhibits two peaks at 6 and 14 eV for the 6 phase and just one peak at 8 eV for the f phase [7]. Palmer and co-workers assigned these 6 and 8 eV peaks to the same ^2P₁ resonant state and attributed the 2-eV shift of the peak energies in the 6 and f phases to multiple electron scattering in the solid [3]. No assignment was made for the 14 eV feature. In this paper, we propose an alternative interpretation for the origin of the 6 and 8 eV peaks and assign the 14 eV peak.

The 6, 8, and 14 eV peak features have been observed also in the ESD of O₂ ions from O₂ films and O₂ on rare gas films, and attributed to the ^2P₁(6eV), ^2P₁(8eV), and ^2P₁(14eV) resonant states, respectively [10]. Excitation to the ^2P₁ and ^2P₃ states from the O₂ ground state is symmetry forbidden in the gas phase but is allowed in the condensed phase when the cylindrical symmetry of the O₂ is relaxed by interaction with the surrounding [11]. Recent ESD studies for O₂ clusters clearly demonstrated that the 8 and 14 eV peaks arise from the "symmetry-forbidden" states [12]. Furthermore, in the gas phase, the symmetries of ^2P₁ and ^2P₃ for the 6 and 8 eV peaks have been established experimentally based on the angular distribution of O₂ ions resulting from electron impact on O₂ in the ground state and the ^3A₁ excited state respectively in line with [13,14]. These assignments are completely in line with theoretical calculations [15] and analysis [11].

2. Origin of the 6, 8, and 14 eV peaks in the X-a excitation
The question here is: Do the 6, 8, and 14 eV peaks observed in the X-a excitation profile arise from the same ^2P₁(6eV),

²P₁(8eV), and ²P₁(14eV) resonant states as those observed in the above experiments? We shall answer this question by examining the angular distributions of inelastically scattered electrons and the electron attachment selectivity observed by Palmer and co-workers [1-8].

2₁P₁(6eV): The 2₁P₁ state was observed at 6 eV in the v=0-1 vibrational excitation profile [7]. The angular distribution of inelastically scattered electrons at incident electron energy E_i=6.5eV for the X-a excitation is almost identical to that for the v=0-1 excitation [7]. This agreement is observed for both the 6 and f phases, suggesting strongly that the 6 eV peaks in the X-a and v=0-1 excitation profiles arise from the same 2₁P₁ state. This may be the most important finding of the present paper.

2₁P₁(8eV): The emitted electrons from the 2₁P₁ state decaying into the a(^1G) state have f symmetry. Therefore, the angular distribution of emitted electrons from this state for the f phase must be sin⁴θ for small θ, where θ is the angle from the direction normal to the graphite surface. This sin⁴θ dependence agrees with experimental results [5]. Further supporting evidence: The 8 eV feature appears in the X-a excitation profile for both the f and 6 phases. (Although there is no discernible peak at 8 eV in the 6 phase, there is a broad shoulder which cannot be accounted for by the 6-eV resonance alone. The 6-eV resonance, which was observed in the v=0-1 excitation, does not have such an extended tail [7].) The 8 eV intensity for the f phase is stronger than that for the 6 phase [7]. This non-selectivity and relative intensity can be explained, if the 8 eV feature arises from the 2₁P₁ resonant state.

2₁P₁(14eV): The angular distribution of the inelastically scattered electrons at 14 eV is consistent with the f symmetry [16]. Further supporting evidence: The 14 eV feature observed in the X-a excitation for the f phase disappears for the 6 phase. Likewise, the ^3P₀(3g) feature observed in the v=0-1 excitation for the 6 phase disappears for the f phase [7]. This parallel behavior is expected, if the 14 eV feature peaks observed in the X-a excitation profile indeed arise from the 2₁P₁ resonant state. Based on the above, we conclude that the 6, 8, and 14 eV arise from the 2₁P₁ resonant state.

Based on the above, we conclude that the 6, 8, and 14 eV

peaks observed in the X-a excitation profile indeed arise from the same 2₁P₁(6eV),

the 2₁P₁(8eV), and 2₁P₁(14eV) states as those

observed in ESD and other experiments. This conclusion disagrees with the assignment made by Palmer and co-workers who attributed the 6 and 8 eV peaks to the 2₁P₁ state.

3. The 2₁P₃ intermediate state in the solid

What happened to the 2₁P₃ state in the solid? In the gas phase, the X-a electronic excitation proceeds via the O₂(^3P₂) intermediate state at least for incident electron energies less than 5 eV [17]. This intermediate state should be differentiated from the O₂(^1S₀) ^3P₂ shape resonance peaked around 0.2 eV [18]. The latter has the 1¹D component predominant within the molecular region and a negligible amplitude outside this region, while the former has comparable amplitudes in both regions. Theoretical calculations [19]

predict that this non-resonant 2^{II}_{Q} intermediate state produces a very broad peak (FWHM=10eV), if there is one, in the X-a excitation function.

The non-resonant 2^{II}_{Q} scattering, unlike the resonant 2^{II}_{Q} scattering, is expected to be perturbed strongly when O_2 is condensed. In fact, there is ample evidence for quenching of this non-resonant 2^{II}_{Q} intermediate state in condensed phase:

- The observed threshold energy (3.0 eV) is delayed by 2.1 eV from the expected threshold energy of 0.9 eV [7,20].
- The angular distribution of inelastically scattered electrons at $E_i=6$ eV indicates 2^{II}_{Q} symmetry rather than 2^{I}_{Q} symmetry, as was argued in Sec. 2.
- The observed features are too sharp (FWHM=4eV) to arise from non-resonant 2^{II}_{Q} scattering. The latter should have either no peak or a very broad peak with FWHM=10 eV. To sum up, the non-resonant 2^{II}_{Q} intermediate state is quenched upon condensation and hence cannot be assigned to the 6 and 8 eV features.

4. Resonance energy shift due to multiple electron scattering

PALMER and co-workers offered two reasons for the 2-eV shift of the 2^{II}_{Q} resonance peaks [3,7]:

- theoretical calculations assuming the 2^{II}_{Q} symmetry and the similarity of the angular distributions for $E_i=6.5$ and 8.5 eV.
- Item (1): The assumption of the 2^{II}_{Q} symmetry is not valid as argued in Sections 2 and 3. This explains partly why their theoretical calculations did not agree well with the experimental data [3,7].
- Item (2): Although the angular distributions for the δ phase, those for the ζ phase are quite different. The only common feature between them is that both decrease toward the normal direction. In short, there is no evidence to support the 2-eV shift of the 2^{II}_{Q} resonance.

5. A concluding remark

PALMER and co-workers assigned the 6 and 8 eV peaks, observed in the X-a excitation, to the 2^{II}_{Q} resonance and attributed the 2-eV shift of the peak energies in the δ and ζ phases to multiple electron scattering in the solid. Based on this single investigation, they claimed that multiple electron scattering can substantially shift resonance energies and warned electron spectroscopists who study adsorbed molecules to be aware of this possibility [3,7]. Moreover, these authors repeatedly suggested this possibility to explain experimental results obtained by others [3,8]. Because of the PALMER and co-workers' claim, other authors considered (although did not adopt) this possibility for an explanation of their results [21]. This paper has demonstrated clearly that the PALMER and co-workers' claim is unfounded at least for the cases studied so far.

Acknowledgment

This work was supported by the Office of Naval Research.

References

- [1] R.E. Palmer, P.J. Rous, J.L. Wilkes and R.F. Willis, Phys. Rev. Lett. 60 (1988) 329.
- [2] P.J. Rous, R.E. Palmer and R.F. Willis, Phys. Rev. B 39 (1989) 7555.
- [3] P.J. Rous, E.T. Jensen and R.E. Palmer, Phys. Rev. Lett. 63 (1989) 2496.
- [4] E.T. Jensen, R.E. Palmer and P.J. Rous, Phys. Rev. Lett. 64 (1990) 1301.
- [5] E.T. Jensen, R.E. Palmer and P.J. Rous, Chem. Phys. Lett. 69 (1990) 204.
- [6] P.J. Rous, R.E. Palmer and E.T. Jensen, Phys. Rev. B 41 (1990) 4793.
- [7] E.T. Jensen, R.E. Palmer and P.J. Rous, Surf. Sci. 237 (1990) 153.
- [8] R.E. Palmer and P.J. Rous, Rev. Modern Phys. April 64 (1992) 383.
- [9] M.F. Tonry and S.C. Pain, Phys. Rev. B 30 (1984) 1115;
- [10] H. You and S.C. Pain, Phys. Rev. B 33 (1986) 5886;
- [11] M.F. Tonry and S.C. Pain, Phys. Rev. B 36 (1987) 1248.
- [12] H. Sambe and D.E. Ramaker, Surf. Sci. (in press).
- [13] H. Sambe and D.E. Ramaker, Phys. Rev. A 40 (1989) 3651.
- [14] R.J. Van Brunt and L.J. Kieffer, Phys. Rev. A 2 (1970) 1899.
- [15] D.S. Bellis and R.I. Hall, J. Phys. B 14 (1981) 365.
- [16] R.H. Michaels, Adv. Chem. Phys. 45 (1981) part 2, 225.
- [17] E.S. Chang, J. Phys. B 10 (1977) L677.
- [18] G.J. Schulz, Rev. Modern Phys. 45 (1973) 423; G.J. Schulz, In Principle of Laser Plasmas edited by G. Bekerfi (John Wiley & Sons, New York, 1976) p.33.
- [19] C.J. Noble and P.G. Burke, J. Phys. B 19 (1986) L35; D. Teillet-Billy, L. Malegat and J.P. Ganyacq, J. Phys. B 20 (1987) 3701.
- [20] L. Sanchez and M. Michaud, Phys. Rev. Lett. 47 (1981) 1008.
- [21] W. Hansen, H. Bertolo and K. Jacob, Surf. Sci. 253 (1991) 1; T.S. Jones, M.R. Ashton, M.Q. Ding, and N.V. Richardson, Chem. Phys. Lett. 161 (1989) 467.

TECHNICAL REPORT DISTRIBUTION LIST - GENERAL

Offices of Naval Research (2)
 Chemistry Division, Code 1113
 800 North Quincy Street
 Arlington, Virginia 22217-5000

Dr. Richard W. Drisko (1)
 Naval Civil Engineering
 Laboratory
 Code 152
 Port Hueneme, CA 93043

Dr. James S. Murday (1)
 Chemistry Division, Code 6100
 Naval Research Laboratory
 Washington, D.C. 20375-5000

Dr. Harold H. Singerman (1)
 Naval Surface Warfare Center
 Carderock Division Detachment
 Annapolis, MD 21402-1198

Dr. Robert Green, Director (1)
 Chemistry Division, Code 385
 Naval Air Weapons Center
 Weapons Division
 China Lake, CA 93555-6001

Dr. Eugene C. Fischer (1)
 Code 2840
 Naval Surface Warfare Center
 Carderock Division Detachment
 Annapolis, MD 21402-1198

Dr. Elik Lindner (1)
 Naval Command, Control and Ocean
 Surveillance Center
 RDT&E Division
 San Diego, CA 92152-5000

Defense Technical Information
 Center (2)
 Building 5, Cameron Station
 Alexandria, VA 22314

Dr. Bernard E. Douda (1)
 Crane Division
 Naval Surface Warfare Center
 Cranes, Indiana 47532-5000

DTIC Selected 1

* Number of copies to forward

Accession For	NTIS GRAFILE	DTIC TAB	Unannounced	Justification
BY-Distribution/	<input checked="" type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>
Availability Codes				
Dist. Special	A-1			